

Mathematical statistics

September 25th, 2018

Lecture 9: Point estimation - Method of moments

Where are we?

Week 1	•	Probability reviews
Week 2	•	Chapter 6: Statistics and Sampling Distributions
Week 4	•	Chapter 7: Point Estimation
Week 7	•	Chapter 8: Confidence Intervals
Week 10	•	Chapter 9: Test of Hypothesis
Week 14	•	Regression

7.1 Point estimate

- unbiased estimator
- mean squared error

7.2 Methods of point estimation

- method of moments
- method of maximum likelihood.


7.3 Sufficient statistic

7.4 Information and Efficiency


- Large sample properties of the maximum likelihood estimator
- Bootstrap

Question of this chapter

- Given a random sample X_1, \dots, X_n from a distribution with pmf/pdf $f(x, \theta)$ parameterized by a parameter θ
- Goal: Estimate θ



Point estimate



Definition

A point estimate $\hat{\theta}$ of a parameter θ is a single number that can be regarded as a sensible value for θ .

$$\begin{array}{ccccc} \text{population parameter} & \implies & \text{sample} & \implies & \text{estimate} \\ \theta & & \implies X_1, X_2, \dots, X_n & \implies & \hat{\theta} \end{array}$$

Mean Squared Error

- Measuring error of estimation

$$|\hat{\theta} - \theta| \quad \text{or} \quad (\hat{\theta} - \theta)^2$$

- The error of estimation is random

Definition

The mean squared error of an estimator $\hat{\theta}$ is

$$E[(\hat{\theta} - \theta)^2]$$


Theorem

$$MSE(\hat{\theta}) = E[(\hat{\theta} - \theta)^2] = V(\hat{\theta}) + \left(E(\hat{\theta}) - \theta\right)^2$$

Bias-variance decomposition

Mean squared error = variance of estimator + (*bias*)²

Bias-variance decomposition



Unbiased estimators

Definition

A point estimator $\hat{\theta}$ is said to be an unbiased estimator of θ if

$$E(\hat{\theta}) = \theta$$

for every possible value of θ .

Unbiased estimator

$$\Leftrightarrow \text{Bias} = 0$$

$$\Leftrightarrow \text{Mean squared error} = \text{variance of estimator}$$

Example 3



Problem

Consider a random sample X_1, \dots, X_n from the pdf

$$f(x) = \frac{1 + \theta x}{2} \quad -1 \leq x \leq 1$$

Show that $\hat{\theta} = 3\bar{X}$ is an unbiased estimator of θ .

Example 7.1 and 7.4



Minimum variance unbiased estimator (MVUE)

Definition

Among all estimators of θ that are unbiased, choose the one that has minimum variance. The resulting $\hat{\theta}$ is called the minimum variance unbiased estimator (MVUE) of θ .

Recall:

- Mean squared error = variance of estimator + $(bias)^2$
- unbiased estimator \Rightarrow bias = 0

\Rightarrow MVUE has minimum mean squared error among unbiased estimators

Estimators of the mean

1.3: Estimators of the mean

- The sample mean
- The sample median
- Trimmed means

Measures of locations: mean

The **sample mean** \bar{x} of observations x_1, x_2, \dots, x_n is given by

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

Measures of locations: median

5, 13, 9, 7, 1, 9, 2, 9, and 11

put in
ascending order

1, 2, 5, 7, 9, 9, 9, 11, 13

Median
(middle value)

Median is not affected by outliers

Measures of locations: median

Ordering the observations from smallest to largest

$$\tilde{x} = \begin{cases} \text{The single middle value if } n \text{ is odd} & = \left(\frac{n+1}{2} \right)^{\text{th}} \text{ ordered value} \\ \text{The average of the two middle values if } n \text{ is even} & = \text{average of } \left(\frac{n}{2} \right)^{\text{th}} \text{ and } \left(\frac{n}{2} + 1 \right)^{\text{th}} \text{ ordered values} \end{cases}$$

Measures of locations: trimmed mean

- A $\alpha\%$ trimmed mean is computed by:
 - eliminating the smallest $\alpha\%$ and the largest $\alpha\%$ of the sample
 - averaging what remains
- $\alpha = 0 \rightarrow$ the mean
- $\alpha \approx 50 \rightarrow$ the median

Problem

Fifteen air samples from a certain region were obtained, and for each one the carbon monoxide concentration was determined. The results (in ppm) were




*19, 21, 17, 20, 24, 30, 18, 23, 19,
26, 22, 17, 27, 25, 22*

- ① *Find the sample mean*
- ② *Find the sample median of the data set*
- ③ *Find the 20% trimmed mean*


What is the best estimator of the mean?

Question: Let X_1, \dots, X_n be a random sample from a distribution with mean μ . What is the best estimator of μ ?

Example 7.8



Normal vs. Cauchy



What is the best estimator of the mean?

Question: Let X_1, \dots, X_n be a random sample from a distribution with mean μ . What is the best estimator of μ ?

Answer: It depends.

- Normal distribution \rightarrow sample mean \bar{X}
- Cauchy distribution \rightarrow sample median \tilde{X}
- Uniform distribution \rightarrow no tails, uniform

$$\hat{X}_e = \frac{\text{largest number} + \text{smaller number}}{2}$$

- In all cases, 10% trimmed mean performs pretty well

MVUE of normal distributions

Theorem

Let X_1, \dots, X_n be a random sample from a normal distribution with mean μ . Then the estimator $\hat{\mu} = \bar{X}$ is the MVUE for μ .

Method of moments

Moments

- Let X_1, \dots, X_n be a random sample from a normal distribution with pmf or pdf $f(x)$.
- For $k = 1, 2, 3, \dots$, the k^{th} population moment, or k^{th} moment of the distribution $f(x)$, is

$$E(X^k)$$

- First moment: the mean
- Second moment: $E(X^2)$

Sample moments

- Let X_1, \dots, X_n be a random sample from a distribution with pmf or pdf $f(x)$.
- For $k = 1, 2, 3, \dots$, the k^{th} sample moment is

$$\frac{X_1^k + X_2^k + \dots + X_n^k}{n}$$

The law of large numbers provides that when $n \rightarrow \infty$

$$\frac{X_1^k + X_2^k + \dots + X_n^k}{n} \rightarrow E(X^k)$$

Method of moments: ideas

- Let X_1, \dots, X_n be a random sample from a distribution with pmf or pdf

$$f(x; \theta_1, \theta_2, \dots, \theta_m)$$

- Assume that for $k = 1, \dots, m$

$$\hat{u}_k = \frac{X_1^k + X_2^k + \dots + X_n^k}{n} = E(X^k)$$

- Solve the system of equations for $\theta_1, \theta_2, \dots, \theta_m$

Method of moments: Example 1

Problem

Let X_1, \dots, X_{10} be a random sample from the exponential distribution with parameter λ , that is

$$f(x; \lambda) = \lambda e^{-\lambda x}, \quad x \geq 0$$

The observed data are

3.11, 0.64, 2.55, 2.20, 5.44,

3.42, 1.39, 8.13, 1.82, 1.30

Use the method of moments to obtain an estimator of λ .

Method of moments: Example 1

- Equation: $k = 1$

$$\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n} = E(X) = \frac{1}{\lambda}$$

- Solve the system of equations for λ

$$\lambda = \frac{1}{\bar{X}}$$

Method of moments: Example 2

Problem

Suppose that for a parameter $0 \leq \theta \leq 1$, X is the outcome of the roll of a four-sided tetrahedral die

x	1	2	3	4
$p(x)$	$\frac{3\theta}{4}$	$\frac{\theta}{4}$	$\frac{3(1-\theta)}{4}$	$\frac{(1-\theta)}{4}$

Suppose the die is rolled 10 times with outcomes

4, 1, 2, 3, 1, 2, 3, 4, 2, 3

Use the method of moments to obtain an estimator of θ .

Method of moments: Example 3

Problem

Let X_1, \dots, X_{10} be a random sample from a distribution with pdf

$$f(x) = \begin{cases} (\theta + 1)x^\theta & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

A random sample of ten students yields data

$$x_1 = .92, x_2 = .79, x_3 = .90, x_4 = .65, x_5 = .86,$$

$$x_6 = .47, x_7 = .73, x_8 = .97, x_9 = .94, x_{10} = .77$$

Use the method of moments to obtain an estimator of θ .

Method of moments: Example 4

Problem

Let X_1, \dots, X_n be a random sample from the normal distribution $\mathcal{N}(0, \sigma^2)$.

Use the method of moments to obtain an estimator of σ .

Method of moments: Example 5

Problem

Let $\beta > 1$ and X_1, \dots, X_n be a random sample from a distribution with pdf

$$f(x) = \begin{cases} \frac{\beta}{x^{\beta+1}} & \text{if } x > 1 \\ 0 & \text{otherwise} \end{cases}$$

Use the method of moments to obtain an estimator of β .