

# MATH 205: Statistical methods

## Lab 8: Propagating uncertainty

# Problem 1

- A farmer wants to know the area of his rectangular field. He asks two probabilists to measure the dimension of the field.
- They did and give him the following summary: Let  $X$ ,  $Y$  be the width and the length of the rectangle, then  $X$  and  $Y$  are independent and

$$X \sim 30 + 3 * \text{Uniform}([0, 1])$$

$$Y \sim 50 + 5 * \text{Beta}(2, 5),$$

- Can you help the farmer find out the mean and the standard deviation of the area?

# Problem 1

- Use two R functions `rnorm` and `rexp` to sample 20000 samples of  $(X, Y)$  and of the area of the field  $A = X \times Y$ .
- Compute the mean, the standard deviation and produce a histogram of  $A$

# Propagation of uncertainty


- In various problem, there is a quantity of interest  $Q$ , modeled as the output of a multivariate function, that is

$$Q = g(X_1, X_2, \dots, X_m)$$

where  $X_1, X_2, \dots, X_m$  are inputs that can be measured (with noise) and  $g$  is a known but complicated function.

- Central question: Assume that we know the distribution of  $X_1, X_2, \dots, X_m$ , can we make prediction about  $Q$ ?
- Answer: Yes
  - Get random samples of  $X_1, X_2, \dots, X_m$
  - Evaluate  $Q = g(X_1, X_2, \dots, X_m) \rightarrow$  obtain samples of  $Q$
  - The histogram of the dataset represent the distribution of  $Q$

## Worldwide nuclear testing, 1945 - 2013



# Black-box models

- In lots of examples, the function  $g$  is so complicated that you don't really know what it does. People refer to such cases as black-box predictions.
- As long as we can evaluate  $g$ , we can sample  $Q$

# Prey-predator model

- Lotka–Volterra equations
- Describes the dynamics of biological systems in which two species interact, one as a predator and the other as prey
- Assume that the two parameters Alpha and Beta in the codes are not constant, but follow the following distributions

$$Alpha, Beta \sim \mathcal{N}(10^{-3}, 10^{-8})$$

- We are interested in  $P$ , the number of preys at the end of the simulation
- Generate 2000 samples of  $P$ . Compute the mean, the standard deviation and produce a histogram of  $P$ .