

# Mathematical statistics

March 6<sup>th</sup>, 2018

## Lecture 10: Point Estimation

# Where are we?

---

<b>Week 1</b> . . . . .	●	Probability reviews
<b>Week 2</b> . . . . .	●	Chapter 6: Statistics and Sampling Distributions
<b>Week 4</b> . . . . .	●	<b>Chapter 7: Point Estimation</b>
<b>Week 7</b> . . . . .	●	Chapter 8: Confidence Intervals
<b>Week 10</b> . . . . .	●	Chapter 9: Test of Hypothesis
<b>Week 14</b> . . . . .	●	Regression

---

## 7.1 Point estimate

- unbiased estimator
- mean squared error

## 7.2 Methods of point estimation

- method of moments
- method of maximum likelihood.


## 7.3 Sufficient statistic


## 7.4 Information and Efficiency

- Large sample properties of the maximum likelihood estimator
- Bootstrap

# Question of this chapter

- Given a random sample  $X_1, \dots, X_n$  from a distribution with pmf/pdf  $f(x, \theta)$  parameterized by a parameter  $\theta$
- Goal: Estimate  $\theta$





## Definition

A point estimate  $\hat{\theta}$  of a parameter  $\theta$  is a single number that can be regarded as a sensible value for  $\theta$ .

population parameter  $\implies$  sample  $\implies$  estimate  
 $\theta \implies X_1, X_2, \dots, X_n \implies \hat{\theta}$

# Example

## Problem

Consider a random sample  $X_1, \dots, X_{10}$  from the pdf

$$f(x) = \frac{1 + \theta x}{2} \quad -1 \leq x \leq 1$$

Assume that the obtained data are

0.92, -0.1, -0.2, 0.75, 0.65, -0.53

0.36, -0.68, 0.97, -0.33, 0.79

Provide an estimate of  $\theta$ .

# Mean Squared Error

- Measuring error of estimation

$$|\hat{\theta} - \theta| \quad \text{or} \quad (\hat{\theta} - \theta)^2$$

- The error of estimation is random

## Definition

The mean squared error of an estimator  $\hat{\theta}$  is

$$E[(\hat{\theta} - \theta)^2]$$

## Problem

Let  $Y$  be a random variable and  $a$  is a constant. Prove that

$$E[(Y - a)^2] = \text{Var}(Y) + (E[Y] - a)^2$$

Hint: Recall that

$$\text{Var}[Y] = E[Y^2] - (E[Y])^2$$




## Theorem

$$MSE(\hat{\theta}) = E[(\hat{\theta} - \theta)^2] = V(\hat{\theta}) + (E(\hat{\theta}) - \theta)^2$$

## Bias-variance decomposition

Mean squared error = variance of estimator + (*bias*)<sup>2</sup>

# Bias-variance decomposition



# Statistical bias vs. social bias

How things should be



## Definition

A point estimator  $\hat{\theta}$  is said to be an unbiased estimator of  $\theta$  if

$$E(\hat{\theta}) = \theta$$

for every possible value of  $\theta$ .

Unbiased estimator

$\Leftrightarrow$  Bias = 0

$\Leftrightarrow$  Mean squared error = variance of estimator

# Sample proportion

- A test is done with probability of success  $p$ . Denote the outcome by let  $X$  (success: 1, failure: 0)

$$E[X] = p, \quad \text{Var}[X] = p(1 - p)$$

- $n$  independent tests are done, let  $X_1, X_2, \dots, X_n$  be the outcomes
- Let

$$\hat{p} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

- We know that

$$E[\hat{p}] = p$$

thus  $\hat{p}$  is an unbiased estimator

- Compute  $MSE(\hat{p})$

- Let

$$\hat{p} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

$$\begin{aligned}MSE(\hat{p}) &= \text{Var}(\hat{p}) + (\text{bias})^2 \\&= \text{Var}\left(\frac{X_1 + X_2 + \dots + X_n}{n}\right) \\&= \frac{1}{n^2} (\text{Var}(X_1) + \text{Var}(X_2) + \dots + \text{Var}(X_n)) \\&= \frac{p(1-p)}{n}\end{aligned}$$

# Sample proportion

- A test is done with probability of success  $p$
- $n$  independent tests are done, let  $X_1, X_2, \dots, X_n$  be the outcomes
- Strange idea: How about using

$$\tilde{p} = \frac{X_1 + X_2 + \dots + X_n + 2}{n + 4}$$

- What is the bias of  $\tilde{p}$ ?
- Compute  $MSE(\tilde{p})$