

Mathematical statistics

March 13th, 2018

Lecture 13: Method of maximum likelihood

Where are we?

Week 1	●	Probability reviews
Week 2	●	Chapter 6: Statistics and Sampling Distributions
Week 4	●	Chapter 7: Point Estimation
Week 7	●	Chapter 8: Confidence Intervals
Week 10	●	Chapter 9: Test of Hypothesis
Week 14	●	Regression

7.1 Point estimate

- unbiased estimator
- mean squared error


7.2 Methods of point estimation

- method of moments
- method of maximum likelihood.

7.3 Sufficient statistic

7.4 Information and Efficiency

- Large sample properties of the maximum likelihood estimator
- Bootstrap



Definition

A point estimate $\hat{\theta}$ of a parameter θ is a single number that can be regarded as a sensible value for θ .

population parameter \implies sample \implies estimate
 $\theta \implies X_1, X_2, \dots, X_n \implies \hat{\theta}$

Method of moments

- The expectation of a exponential random variable is

$$E[X] = \frac{1}{\lambda}$$

- For large n , we have

$$\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

is close to $E[X]$

- We can compute \bar{x} from the data \rightarrow approximate λ

- Let X_1, \dots, X_n be a random sample from a normal distribution with pmf or pdf $f(x)$.
- For $k = 1, 2, 3, \dots$, the k^{th} population moment, or k^{th} moment of the distribution $f(x)$, is

$$E(X^k)$$

- First moment: the mean
- Second moment: $E(X^2)$

Sample moments

- Let X_1, \dots, X_n be a random sample from a distribution with pmf or pdf $f(x)$.
- For $k = 1, 2, 3, \dots$, the k^{th} sample moment is

$$\frac{X_1^k + X_2^k + \dots + X_n^k}{n}$$

The law of large numbers provides that when $n \rightarrow \infty$

$$\frac{X_1^k + X_2^k + \dots + X_n^k}{n} \rightarrow E(X^k)$$

- Let X_1, \dots, X_n be a random sample from a distribution with pmf or pdf

$$f(x; \theta_1, \theta_2, \dots, \theta_m)$$

- Assume that for $k = 1, \dots, m$

$$\hat{u}_k = \frac{X_1^k + X_2^k + \dots + X_n^k}{n} = E(X^k)$$

- Solve the system of equations for $\theta_1, \theta_2, \dots, \theta_m$

Method of moments: Example 2

Problem

Suppose that for a parameter $0 \leq \theta \leq 1$, X is the outcome of the roll of a four-sided tetrahedral die

x	1	2	3	4
$p(x)$	$\frac{3\theta}{4}$	$\frac{\theta}{4}$	$\frac{3(1-\theta)}{4}$	$\frac{(1-\theta)}{4}$

Suppose the die is rolled 10 times with outcomes

4, 1, 2, 3, 1, 2, 3, 4, 2, 3

Use the method of moments to obtain an estimator of θ .

Problem

Let X_1, \dots, X_{10} be a random sample from a distribution with pdf

$$f(x) = \begin{cases} (\theta + 1)x^\theta & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

A random sample of ten students yields data

$$x_1 = .92, x_2 = .79, x_3 = .90, x_4 = .65, x_5 = .86,$$

$$x_6 = .47, x_7 = .73, x_8 = .97, x_9 = .94, x_{10} = .77$$

Use the method of moments to obtain an estimator of θ .

Problem

Let X_1, \dots, X_n be a random sample from the normal distribution $\mathcal{N}(0, \sigma^2)$.

Use the method of moments to obtain an estimator of σ .


Problem

Let $\beta > 1$ and X_1, \dots, X_n be a random sample from a distribution with pdf

$$f(x) = \begin{cases} \frac{\beta}{x^{\beta+1}} & \text{if } x > 1 \\ 0 & \text{otherwise} \end{cases}$$

Use the method of moments to obtain an estimator of β .

Method of maximum likelihood



Definition

The random variables X_1, X_2, \dots, X_n are said to form a (simple) random sample of size n if

- 1 the X_i 's are independent random variables
- 2 every X_i has the same probability distribution

Definition

Two random variables X and Y are said to be independent if for every pair of x and y values,

$$P(X = x, Y = y) = P_X(x) \cdot P_Y(y) \quad \text{if the variables are discrete}$$

or

$$f(x, y) = f_X(x) \cdot f_Y(y) \quad \text{if the variables are continuous}$$

Let X_1, X_2, \dots, X_n be a random sample of size n from a distribution with density function $f_X(x)$.

Then the density of the joint distribution of (X_1, X_2, \dots, X_n) is

$$f_{joint}(x_1, x_2, \dots, x_n) = \prod_{i=1}^n f_X(x_i)$$

Maximum likelihood estimator

- Let X_1, X_2, \dots, X_n have joint pmf or pdf

$$f_{joint}(x_1, x_2, \dots, x_n; \theta)$$

where θ is unknown.

- When x_1, \dots, x_n are the observed sample values and this expression is regarded as a function of θ , it is called the likelihood function.
- The maximum likelihood estimates θ_{ML} are the value for θ that maximize the likelihood function:

$$f_{joint}(x_1, x_2, \dots, x_n; \theta_{ML}) \geq f_{joint}(x_1, x_2, \dots, x_n; \theta) \quad \forall \theta$$

How to find the MLE?

- Step 1: Write down the likelihood function.
- Step 2: Can you find the maximum of this function?
- Step 3: Try taking the logarithm of this function.
- Step 4: Find the maximum of this new function.

To find the maximum of a function of θ :

- compute the derivative of the function with respect to θ
- set this expression of the derivative to 0
- solve the equation

Problem

Let X_1, \dots, X_{10} be a random sample from the exponential distribution with parameter λ , that is

$$f(x; \lambda) = \lambda e^{-\lambda x}, \quad x \geq 0$$

The observed data are

3.11, 0.64, 2.55, 2.20, 5.44,

3.42, 1.39, 8.13, 1.82, 1.30

Use the method of maximum likelihood to obtain an estimator of λ .

Example

Problem

Let X_1, \dots, X_{10} be a random sample from a distribution with pdf

$$f(x) = \begin{cases} (\theta + 1)x^\theta & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

A random sample of ten students yields data

$$x_1 = .92, x_2 = .79, x_3 = .90, x_4 = .65, x_5 = .86,$$

$$x_6 = .47, x_7 = .73, x_8 = .97, x_9 = .94, x_{10} = .77$$

Use the method of maximum likelihood to obtain an estimator of θ .

Example

Problem

Let X_1, \dots, X_n be a random sample from the normal distribution $\mathcal{N}(0, \sigma^2)$, that is

$$f(x, \theta) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{x^2}{2\sigma^2}}$$

Use the method of maximum likelihood to obtain an estimator of σ .

Example

Problem

Let $\beta > 1$ and X_1, \dots, X_n be a random sample from a distribution with pdf

$$f(x) = \begin{cases} \frac{\beta}{x^{\beta+1}} & \text{if } x > 1 \\ 0 & \text{otherwise} \end{cases}$$

Use the method of maximum likelihood to obtain an estimator of β .