

Mathematical techniques in data science


Lecture 30: Metropolis-Hasting algorithm

May 3rd, 2019

Schedule

Week	Chapter
1	Chapter 2: Intro to statistical learning
3	Chapter 4: Classification
4	Chapter 9: Support vector machine and kernels
5, 6	Chapter 3: Linear regression
7	Chapter 8: Tree-based methods + Random forest
8	
9	Neural networks
12	PCA → Manifold learning
11	Clustering: K-means → Spectral Clustering
10	Bayesian methods + UQ
13	Reinforcement learning/Online learning/Active learning
14	Project presentation

Uncertainty quantification



More generally: suppose we have a model for X that depends on some parameters θ . Then:

- 1 Choose a prior $P(\theta)$ for θ .
- 2 Compute the posterior distribution of θ using

$$p(\theta|X) \propto P(X|\theta) \cdot P(\theta).$$

Note: Posterior = Prior \times Likelihood.

How do we sample from the posterior distribution

$$P(\theta | data) \propto P(data|\theta).P(\theta)$$

assuming that we can evaluate $P(\theta | data)$ point-wise (up to a normalizing constant)

Markov chain Monte Carlo

Markov chains

- Let $S := \{s_1, s_2, \dots\}$ be a countable set.
- A (discrete time) **Markov chain** is a discrete stochastic process $\{X_n : n = 0, 1, \dots\}$ such that
 - ① X_n is an S -valued random variable $\forall n \geq 0$.
 - ② (Markov Property) For all $i, j, i_0, \dots, i_{n-1} \in S$, and all $n \geq 0$:
$$P(X_{n+1} = j | X_0 = i_0, \dots, X_{n-1} = i_{n-1}, X_n = i) = P(X_{n+1} = j | X_n = i).$$

Interpretation: Given the present X_n , the future X_{n+1} is independent of the past (X_0, \dots, X_{n-1}) .

- The elements of S are called the **states** of the Markov chain.
- When $X_n = j$, we say that the process is in state j at time n .

Stationary Markov chains

- A Markov chain is **homogeneous** (or **stationary**) if for all $n \geq 0$ and all $i, j \in S$,

$$P(X_{n+1} = j | X_n = i) = P(X_1 = j | X_0 = i) =: p(i, j).$$

In other words, the **transition probabilities** do not depend on time.

- We will only consider homogeneous chains in what follows.

Transition probabilities

- We denote by $P := (p(i, j))_{i, j \in S}$ the **transition matrix** of the chain.
- Note: P is a **stochastic matrix**, i.e.,

$$\forall i, j \in S, p(i, j) \geq 0, \quad \text{and} \quad \forall i \in S, \sum_{j \in S} p(i, j) = 1.$$

- Conversely, every stochastic matrix is the transition matrix of some homogeneous discrete time Markov chain.


Example

Example 1: (Two-state Markov chain)

$$S = \{0, 1\}, \quad p(0, 1) = a, \quad p(1, 0) = b, \quad a, b \in [0, 1]$$

$$P = \begin{pmatrix} 1-a & a \\ b & 1-b \end{pmatrix}.$$

We naturally represent P using a transition (or state) diagram:



Let $\{X_n : n \geq 0\}$ be a Markov chain.

- We define the **initial distribution** of the chain by

$$\mu_0(i) := P(X_0 = i) \quad (i \in S).$$

- All distributional properties of a (homogeneous) Markov Chain are determined by its initial distribution and transition probability matrix.
- For $n \geq 1$, we define the **n -step transition probability** $p^n(i, j)$ by

$$p^n(i, j) := P(X_n = j | X_0 = i) = P(X_{n+m} = j | X_m = i).$$

Also, define

$$p^0(i, j) = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}.$$

- We define the **n -step transition matrix** by

$$P^{(n)} := (p^n(i, j) : i, j \in S).$$

Theorem: (The Chapman-Kolmogorov Equations) We have for all $m, n \geq 1$:

$$P^{(n+m)} = P^{(n)} \cdot P^{(m)}.$$

In particular, for all $n \geq 1$,

$$P^{(n)} = P \cdot P^{(n-1)} = \dots = P^n.$$

Moral: n -step transition probabilities are computed using matrix multiplications.

- Let $\mu_n := (\mu_n(i) : i \in S)$ denote the **distribution of X_n** :

$$\mu_n(i) := P(X_n = i).$$

Proposition: We have

$$\mu_{m+n} = \mu_m P^n, \quad \text{and} \quad \mu_n = \mu_0 P^n.$$

Moral: Distributional computations for Markov Chains are just matrix multiplications.

- **Reducibility:**

- A state $j \in S$ is said to be **accessible** from $i \in S$ (denoted $i \rightarrow j$) if a system started in state i has a non-zero probability of transitioning into state j at some point.
- A state $i \in S$ is said to **communicate** with state $j \in S$ (denoted $i \leftrightarrow j$) if both $i \rightarrow j$ and $j \rightarrow i$.

Note: Communication is an equivalence relation.

A Markov chain is said to be **irreducible** if its state space is a single communicating class.

- **Transience:**

- A state $i \in S$ is said to be **transient** if, given that we start in state i , there is a non-zero probability that we will never return to i .
- A state is **recurrent** if it is not transient.
- The **recurrence time** of state $i \in S$ is $T_i := \min\{n \geq 1 : X_n = i \text{ given } X_0 = i\}$.
- Note: $i \in S$ is recurrent iff $P(T_i < \infty) = 1$.

- **Periodicity:**

- A state $i \in S$ has period k if

$$k = \gcd\{n > 0 : P(X_n = i | X_0 = i) > 0\}.$$

For example, suppose you start in state i and can only return to i at time 6, 8, 10, 12, etc.. Then the period of i is 2.

- If $k = 1$, then the state is said to be aperiodic.

A Markov chain is **aperiodic** if every state is aperiodic.

Limiting behavior of Markov chains: What happens to $p^n(i, j)$ as $n \rightarrow \infty$?

Example: (The two-state Markov chain)



Limiting behavior

Recalling that

$$S = \{0, 1\}, \quad p(0, 1) = a, \quad p(1, 0) = b, \quad a, b \in [0, 1]$$

$$P = \begin{pmatrix} 1-a & a \\ b & 1-b \end{pmatrix}.$$

If $(a, b) \neq (0, 0)$, we have (exercise):

$$P^n = \frac{1}{a+b} \begin{pmatrix} b & a \\ b & a \end{pmatrix} + \frac{(1-a-b)^n}{a+b} \begin{pmatrix} a & -a \\ -b & b \end{pmatrix}.$$

Thus, if $(a, b) \neq (0, 0)$ and $(a, b) \neq (1, 1)$, then

$$\begin{aligned} \lim_{n \rightarrow \infty} p^n(0, 0) &= \lim_{n \rightarrow \infty} p^n(1, 0) = \frac{b}{a+b} \\ \lim_{n \rightarrow \infty} p^n(0, 1) &= \lim_{n \rightarrow \infty} p^n(1, 1) = \frac{a}{a+b}. \end{aligned}$$

Limiting behavior

$$\begin{aligned}\lim_{n \rightarrow \infty} p^n(0, 0) &= \lim_{n \rightarrow \infty} p^n(1, 0) = \frac{b}{a+b} \\ \lim_{n \rightarrow \infty} p^n(0, 1) &= \lim_{n \rightarrow \infty} p^n(1, 1) = \frac{a}{a+b}.\end{aligned}$$

Thus, the chain has a **limiting distribution**.

The limiting distribution is **independent of the initial state**.

Stationary distribution

Recall: $\mu_{n+1} = \mu_n P$.

A vector $\pi = (\pi(i) : i \in S)$ is said to be a **stationary distribution** for a Markov chain $\{X_n : n \geq 0\}$ if

- 1 $0 \leq \pi_i \leq 1 \forall i \in S$.
- 2 $\sum_{i \in S} \pi_i = 1$.
- 3 $\pi = \pi P$, where P is the transition probability matrix of the Markov chain.

Stationary distribution

Remark: In general, a stationary distribution may not exist or be unique.

Theorem: Let $\{X_n : n \geq 0\}$ be an irreducible and aperiodic Markov chain where each state is positive recurrent. Then

- 1 The chain has a unique stationary distribution π .
- 2 For all $i \in S$, $\lim_{n \rightarrow \infty} P(X_n = i) = \pi(i)$.
- 3 $\pi_i = \frac{1}{E[T_i]}$.

$\pi(i)$ can be interpreted as the average proportion of time spent by the chain in state i .

Detailed balance condition

- A Markov chain is said to satisfy the *detailed balance* condition with respect to a probability distribution π if

$$\pi_i P_{ij} = \pi_j P_{ji}$$

for all states i and j .

- Detailed balance \Rightarrow stationary

$$[\pi P]_i = \sum_j \pi_j P_{ji} = \sum_j \pi_i P_{ij} = \pi_i \sum_j P_{ij} = \pi_i$$

- Classical Markov chain theory: given a transition matrix (transition probability), prove that the chain converges to some stationary distribution
- Markov chain Monte Carlo: given a desired distribution π , construct a transition probability such that a chain with that probabilistic law converges to π

- Markov chain Monte Carlo (MCMC) methods are popular ways of sampling from complicated distributions (e.g. the posterior distribution of a complicated model).
- Idea:
 - 1 Construct a **Markov chain** with the desired distribution as its **stationary distribution** π .
 - 2 Burn (e.g. forget) a given number of samples from the Markov chain (while the chain converges to its stationary distribution).
 - 3 Generate a sample from the desired distribution (approximately).

Metropolis-Hastings algorithm

- Nicolas Metropolis (1915–1999) was an American physicist. He worked on the first nuclear reactors at the Los Alamos National Laboratory during the second world war. Introduced the algorithm in 1953 in the paper

Equation of State Calculations by Fast Computing Machines

with A. Rosenbluth, M. Rosenbluth, A. Teller, and E. Teller

- W. K. Hastings (Born 1930) is a Canadian statistician who extended the algorithm to the more general case in 1970.

Metropolis-Hastings algorithm

- Suppose we want to sample from a distribution $P(x) = f(x)/K$, where $K > 0$ is some constant.

Note: The normalization constant K is often unknown and difficult to compute.

- The Metropolis–Hastings starts with an initial sample, and generate new samples using a *transition probability density* $q(x, y)$ (the *proposal distribution*).
- We assume
 - we can evaluate $f(x)$ at every x .
 - we can evaluate $q(x, y)$ at every x, y .
 - we can sample from the distribution $q(x, \cdot)$.

Metropolis-Hastings algorithm

The Metropolis–Hastings algorithm: we start with x_0 such that $f(x_0) > 0$. For $i = 0, \dots$

- 1 Generate a new value y according to $q(x, \cdot)$.
- 2 Compute the “Hastings” ratio:

$$R = \frac{f(y)q(y, x)}{f(x)q(x, y)}$$

- 3 “Accept” the new sample y with probability $\min(1, R)$. If y is accepted, set $x_{i+1} := y$. Otherwise, $x_{i+1} = x_i$.

$$\begin{aligned}f(x)P(x, y) &= f(x)q(x, y) \cdot \min\left(1, \frac{f(y)q(y, x)}{f(x)q(x, y)}\right) \\&= \min(f(x)q(x, y), f(y)q(y, x)) \\&= f(y)q(y, x) \cdot \min\left(\frac{f(x)q(x, y)}{f(y)q(y, x)}, 1\right) \\&= f(y)P(y, x)\end{aligned}$$

Metropolis-Hastings algorithm

Some difficulties:

- Choosing an efficient proposal distribution $q(x, y)$.
- How long should we wait for the Markov chain to converge to the desired distribution, i.e., how many samples should we burn?
- How long should we sample after convergence to make sure we sample in low probability regions?

Gibbs sampling

- Idea: use the conditional distribution of X to generate new samples.
- Note: only possible when the conditional distributions are “nice”. Suppose $X = (X_1, \dots, X_p)$ and $X^{(i)} = (x_1^{(i)}, \dots, x_p^{(i)})$ is a given sample. Generate a new sample $X^{(i+1)} = (x_1^{(i+1)}, \dots, x_p^{(i+1)})$ as follows:

- 1 Generate $x_1^{(i+1)}$ according to the marginal

$$p(x_1 | x_2^{(i)}, \dots, x_p^{(i)}).$$

- 2 Generate $x_2^{(i+1)}$ according to

$$p(x_2 | x_1^{(i+1)}, x_3^{(i)}, \dots, x_p^{(i)}).$$

- 3 Generate $x_3^{(i+1)}$ according to

$$p(x_3 | x_1^{(i+1)}, x_2^{(i+1)}, x_4^{(i)}, \dots, x_p^{(i)}).$$